

UNCLASSIFIED

AD NUMBER
ADB254366
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies only; Proprietary Info.; Mar 2000. Other requests shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott St., Fort Detrick, MD 21702-5012.
AUTHORITY
USAMRMC ltr, 19 Jan 2001.

THIS PAGE IS UNCLASSIFIED

AD _____

Award Number: DAMD17-98-1-8010

TITLE: A Comparison of Post-Deployment Hospitalization
Incidence Between Vietnam and Gulf War Veterans

PRINCIPAL INVESTIGATOR: Christopher G. Blood, Ph.D.
Tabatha L. Aboumrads

CONTRACTING ORGANIZATION: Naval Health Research Center
San Diego, California 92186-5122

REPORT DATE: March 2000

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel
Command, Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Distribution authorized to U.S. Government
agencies only (proprietary information, Mar 00). Other requests
for this document shall be referred to U.S. Army Medical Research
and Materiel Command, 504 Scott Street, Fort Detrick, Maryland
21702-5012.

The views, opinions and/or findings contained in this
report are those of the author (s) and should not be
construed as an official Department of the Army position,
policy or decision unless so designated by other
documentation.

20000607 088

NOTICE

USING GOVERNMENT DRAWINGS, SPECIFICATIONS, OR OTHER DATA INCLUDED IN THIS DOCUMENT FOR ANY PURPOSE OTHER THAN GOVERNMENT PROCUREMENT DOES NOT IN ANY WAY OBLIGATE THE U.S. GOVERNMENT. THE FACT THAT THE GOVERNMENT FORMULATED OR SUPPLIED THE DRAWINGS, SPECIFICATIONS, OR OTHER DATA DOES NOT LICENSE THE HOLDER OR ANY OTHER PERSON OR CORPORATION; OR CONVEY ANY RIGHTS OR PERMISSION TO MANUFACTURE, USE, OR SELL ANY PATENTED INVENTION THAT MAY RELATE TO THEM.

LIMITED RIGHTS LEGEND

Award Number: DAMD17-98-1-8010

Organization: Naval Health Research Center

Location of Limited Rights Data (Pages):

Those portions of the technical data contained in this report marked as limited rights data shall not, without the written permission of the above contractor, be (a) released or disclosed outside the government, (b) used by the Government for manufacture or, in the case of computer software documentation, for preparing the same or similar computer software, or (c) used by a party other than the Government, except that the Government may release or disclose technical data to persons outside the Government, or permit the use of technical data by such persons, if (i) such release, disclosure, or use is necessary for emergency repair or overhaul or (ii) is a release or disclosure of technical data (other than detailed manufacturing or process data) to, or use of such data by, a foreign government that is in the interest of the Government and is required for evaluational or informational purposes, provided in either case that such release, disclosure or use is made subject to a prohibition that the person to whom the data is released or disclosed may not further use, release or disclose such data, and the contractor or subcontractor or subcontractor asserting the restriction is notified of such release, disclosure or use. This legend, together with the indications of the portions of this data which are subject to such limitations, shall be included on any reproduction hereof which includes any part of the portions subject to such limitations.

THIS TECHNICAL REPORT HAS BEEN REVIEWED AND IS APPROVED FOR PUBLICATION.

LTC Karl Friedl

REPORT DOCUMENTATION PAGEForm Approval
OMD No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for receiving instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA. 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 2000		3. REPORT TYPE AND DATE COVERED Final (23 Feb 98 -22 Feb 00)	
4. TITLE AND SUBTITLE A Comparison of Post-Deployment Hospitalization Incidents Between Vietnam and Gulf War Veterans				5. FUNDING NUMBERS DAMD17-98-1-8010	
6. AUTHOR(S) Christopher G. Blood, Ph.D. Tabatha L. Aboumrads					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center San Diego, California 92186-5122				8. PERFORMING ORGANIZATION	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT Distribution authorized to U.S. Government agencies only (proprietary information, Mar 00). Other requests for this document shall be referred to U.S. Army Medical Research and Materiel Command, 504 Scott Street, Fort Detrick, Maryland 21702-5012.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Post-conflict hospital admission rates of Marine Corps units deployed to the Gulf War were compared with the rates of similar units returning from the Vietnam Conflict. Overall, and in four of the five postdeployment years, the aggregated Gulf War veterans exhibited lower hospitalization rates than did their Vietnam counterparts. Separately, the infantry and service support units deployed to the Gulf War theater had lower postdeployment hospitalization rates than did the Vietnam veterans. There were no significant hospitalization rate differences between the cohorts of combat engineer and artillery units. Of the veterans that were hospitalized, personnel from both conflicts averaged 1.24 admissions. In comparing hospital admission categories, differences existed between the Vietnam and Gulf War veterans in their proportions of 'Infective and Parasitic' diseases, 'Genitourinary' disorders, and 'Musculoskeletal' disorders; Vietnam veterans had the higher proportions within the first two categories, and Gulf War veterans had a higher proportion of musculoskeletal disorders. The data suggest that Gulf War veterans are not at an increased risk of hospitalization when contrasted with veterans returning from a previous combat engagement, and that the categories of illnesses for which Gulf War veterans are hospitalized do not substantially differ from those incurred by a previously-deployed combat cohort.					
14. SUBJECT TERMS Marines, Gulf War, postdeployment illness, hospitalizations				15. NUMBER OF PAGES 4	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified		19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	
20. LIMITATION OF ABSTRACT Unlimited					

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

 Where copyrighted material is quoted, permission has been obtained to use such material.

 Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

 Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

N/A In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and use of Laboratory Animals of the Institute of Laboratory Resources, national Research Council (NIH Publication No. 86-23, Revised 1985).

I For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

N/A In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

N/A In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

N/A In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Christopher J. Block 3/15/00

PI - Signature Date

Table of Contents

(1)	Cover	
(2)	SF 298	2
(3)	Foreword	3
(4)	Table of Contents	4
(5)	Introduction	5
(6)	Body	5
(7)	Key Research Accomplishments	6
(8)	Reportable Outcomes	6
(9)	Conclusions	6
(10)	References	7
(11)	Bibliography of Publications and Meeting Abstracts	7
(12)	List of Personnel	7
(13)	Appendices	8

INTRODUCTION

A wide range of postdeployment health problems have been reported among Gulf War veterans. Often these problems are attributed to some yet-undefined environmental exposure that occurred during the veterans' deployment to the Gulf War theater of operations. The present study sought to investigate any differences or similarities between the post-war hospitalization experiences of Marines deployed to the Gulf War and those of a similar group of Marines deployed to an earlier combat engagement – the Vietnam Conflict. This investigation explored two different aspects of hospitalization incidence among the two cohorts. First, the study contrasted the postdeployment hospital admission rates of Marines serving in the two conflicts. Second, the various illness/injury categories of the hospital admissions were contrasted for the two groups of veterans. Hospitalization incidence was further analyzed by four types of troops deployed to both conflicts: infantry, artillery, combat engineer, and service support.

BODY

The statement of work outlined specific tasks necessary in order to perform the study:

- obtain Unit Diaries of Marine combat, combat support, and service support units deployed to Vietnam;
- examine Unit Diaries and any other relevant resources to determine combat activity levels of specific units;
- extract personnel identifiers of Marines in the Vietnam-deployed units and compile into data bases;
- extract personnel identifiers for selected Marine Gulf War-deployed units;
- determine activities of selected Gulf War units from all available sources;
- extract service and medical history data on study populations and compile into data bases
- define criteria for a single hospitalization
- determine total number of person-days within each of the combat, combat support, and service support groups by examining post-deployment lengths of service;
- compute rates of hospital admissions for each group by examining overall admissions and specific diagnostic category admissions;
- contrast and compare for statistical significance any rate differences between the study populations.
- Compile findings into technical report

All of the above tasks were accomplished and are documented in the report, *A Comparison of Postdeployment Hospitalization Incidence Between Vietnam and Gulf War Veterans*, which is included as Appendix A. One task – Compute incidence of medical disability discharge – could not be performed because of data unavailability for the Vietnam cohort.

KEY RESEARCH ACCOMPLISHMENTS

- Report compiled which compares and contrasts the postdeployment hospitalization experiences of Gulf War and Vietnam veterans.

REPORTABLE OUTCOME

The manuscript has been submitted to the American Journal of Epidemiology for publication consideration and will be published as a technical report by the local Defense Automated Printing Service.

CONCLUSIONS

The present study sought to investigate any differences or similarities between the post-war hospitalization experiences of Gulf War and Vietnam veterans. The overall hospital admission rates of the Gulf War and Vietnam veterans in this study are based on 9.6 million and 6.6 million postdeployment person-days respectively, and therefore provide considerable insights into the post-conflict experiences of two groups of veterans. The overall hospital admission rate of the Gulf War veterans of this study was significantly lower than the rate of troops deployed to the last ground conflict with sustained US involvement. Examined individually, the infantry and service support units deployed to the Gulf exhibited significantly lower hospitalization rates following the conflict than did their Vietnam counterparts; the rates of the engineer and artillery units deployed to the Gulf War did not differ significantly from similar units deployed to Vietnam.

Vietnam veterans had significantly higher percentages of their overall admissions accounted for by infective/parasitic disorders and genitourinary disorders than did the Gulf War veterans. In contrast, among Gulf War-deployed Marines the musculoskeletal disorder category represented a significantly higher proportion of their overall admissions than was observed among the Vietnam veterans. The types of musculoskeletal admissions incurred by Gulf War veterans, however, did not differ substantially from the types of musculoskeletal disorders incurred by their Vietnam counterparts. With the exception of these categories of hospital admissions, illness category percentages between the Gulf War and Vietnam veterans exhibited but minimal differences. Among

individuals who were hospitalized, the average number of admissions was the same for the Gulf War and Vietnam cohorts.

The lack of differences between the hospitalization rates of various Gulf War units (infantry, artillery, engineer, service support) performing different functions in varying locations on the battlefield, undercuts the notion that a specific environmental exposure is responsible for deleterious health effects on Gulf War veterans; any specific risk factor might have been expected to affect these very different groups disproportionately. Indeed, Marines deployed to the Gulf War theater exhibited rates of hospitalization that were lower or comparable to the rates incurred by Marines returning from a previous ground combat operation. Nevertheless, the potential psychological disruption associated with deploying a young man or woman across the globe to risk his or her life defending national or international interests, under sometimes harrowing battlefield conditions, is not insubstantial. All appropriate measures should be taken to adequately prepare such individuals before deployment, to monitor their mental and physical well-being during the deployment, and to provide any appropriate counseling or treatment following their return from war.

REFERENCES

References are included on pages 14 to 16 of the appended manuscript.

BIBLIOGRAPHY OF PUBLICATIONS AND MEETING ABSTRACTS

This section is not applicable to the current study.

LIST OF PERSONNEL RECEIVING PAY FROM THE RESEARCH EFFORT

Christopher Blood, M.A.
Tabatha Aboumrad, M.P.H.
Cathy Lam, B.A.
Jeannine Lane, Ph.D.
Elizabeth Brehm, B.A.
Sandy Chu, B.A.
Jay Walker, B.S.

APPENDICES

Appendix A is the final manuscript, *A Comparison of Postdeployment Hospitalization Incidence Between Vietnam and Gulf War Veterans*. Appendix B is the Principal Investigator's curriculum vitae.

APPENDIX A

**A COMPARISON OF POSTDEPLOYMENT HOSPITALIZATION INCIDENCE
BETWEEN VIETNAM AND GULF WAR VETERANS**

Christopher G. Blood, M.A.
Tabatha L. Aboumrads, MPH

Medical Information Systems and
Operations Research Department

Naval Health Research Center
P.O. Box 85122
San Diego, CA 92186-5122

Naval Health Research Center Report No. 00-02 , supported by the U.S. Army Medical Research Acquisition Activity, Fort Detrick, Maryland under Grant No. DAMD17-98-8010. The views expressed in this article are those of the authors and do not reflect the official policy of the Department of Defense, nor the U.S. Government. Approved for public release; distribution unlimited.

The authors gratefully acknowledge the following individuals: Paula Konoske, Ph.D. for her input to the design of this study; Greg Gray, M.D., MPH for his help in study design and manuscript review; Jeannine Lane, Ph.D. for her contributions to the development of the Vietnam database; Tyler Smith, MS for performing data extractions; and Cathy Lam for data management.

A Comparison of Postdeployment Hospitalization Incidence Between Vietnam and Gulf War Veterans

Accounts of postdeployment health problems incurred by Gulf War veterans are many and varied. These accounts have reported a wide range of symptomatology, which have included fatigue, sleep disturbances, joint pain, muscle weakness, depression, inability to concentrate, dental anomalies, skin rashes, gastrointestinal disturbances, headaches, and other health problems.¹⁻⁸ Paralleling the numerous maladies attributed to Gulf War deployment are the many causes advanced as responsible for the medical irregularities. The causes put forth include exposures to chemical/nerve agents,⁹⁻¹¹ exposure to petrochemicals associated with the Kuwait oil fires,¹² exposure to depleted uranium rounds on the battlefield,¹³ ingestion of pyridostigmine bromide,¹⁴ contraction of an infectious disorder such as leishmaniasis,¹⁵ multiple chemical sensitivities,^{16,17} and combat stress.¹⁸⁻²⁰ In short, many Gulf War veterans report symptomatology that they attribute to their deployment to the Gulf War theater of operations.

However, large epidemiologic studies^{21,22} have indicated that postdeployment hospitalizations and deaths among Gulf War veterans have been incurred at rates not statistically different than their nondeployed counterparts. These studies, examining more than one-half million soldiers, sailors, marines, and airmen deployed to the Gulf War theater of operations between August 1990 and mid-1991, and following them for two years after their return from the deployment, provide support to the notion that if a causal link does exist between a deployment-related exposure and ensuing medical anomalies, such a link is not manifested uniformly throughout the entire population of deployed individuals.

The present investigation seeks to shed light on two different aspects of hospitalization incidence among Gulf War Veterans (GWVs). First, while similarities in the 'syndromes' afflicting substantial numbers of veterans returning from various combat engagements have been documented,²³ a systematic contrast of postdeployment hospitalization rates from differing conflicts has not yet been conducted. The present

study compares the postdeployment hospitalization incidence of US Marine Corps (USMC) veterans returning from two different combat engagements – the Vietnam Conflict and the Gulf War. Second, any environmental exposures during the Gulf War having long-term deleterious health effects would be expected to disproportionately affect troops according to their duties or deployment locations – it is unlikely that all personnel deployed to a large theater of operations would have the same rate of exposure to a specific noxious agent. In an attempt to gain insights into the effects of potentially different exposures, the present study further analyzed the hospitalization incidence of Gulf War and Vietnam veterans by contrasting the admission rates for varying troop categories (infantry troops, artillery troops, combat engineers, and service support troops) that had differing duties and battlefield locations during their combat deployments.

METHODS

Data Sources – Personnel

Marine Corps units returning from Vietnam at the end of USMC involvement in that conflict (March – June, 1971 time frame) were first identified via a Marine Corps history,²⁴ and then diaries for these units were obtained from the Records Service Section, Headquarters USMC. The units returning in this time frame were all 1st Marine Division troops and included six infantry battalions (the 1st, 2nd, and 3rd battalions of both the 1st and 5th Marine Regiments); two artillery battalions (1st and 2nd battalions of the 11th Artillery Regiment); one engineer battalion (1st Engineer Battalion); and four service support battalions from the 1st Force Service Regiment (Headquarters and Service Battalion, Supply Battalion, Motor Transport Battalion, and Maintenance Battalion).

Diaries for these units identified 16,980 unique Service Numbers (SNs). Because Marine Corps medical records starting in the 1970s identify individuals by social security number (SSN) rather than SN, the service numbers from the unit diaries were then matched against a file maintained at the Defense Manpower Data Center (DMDC), Monterey Bay, California, that contained both SNs and their corresponding SSNs. Of the

16,980 SNs, social security numbers were ascertainable for 15,194 personnel (89.5%). Nonmatches were not systematically tied to any particular units but instead likely resulted from incomplete data as the military services transitioned from using SNs to SSNs at a time when many individuals were exiting the service.

The SNs and SSNs were then matched against Department of Defense (DoD) Loss Files, also maintained by DMDC, that indicated the date each individual separated from service. Of the 15,194 personnel whose SSN and SN were both ascertainable, 3300 were excluded from the study for the following reasons: 81 personnel were either officers or were members of another service branch whose medical histories were unavailable for this study; 1637 personnel showed a separation from service immediately upon return from Vietnam and therefore could not be medically tracked for any period of time; and service departure date was indeterminable on 1582 individuals.

Thus, 11,894 enlisted male Marines were identified who served in Vietnam, and who did not separate from military service immediately upon their return from South East Asia. These 11,894 men were comprised of 7009 Marines in infantry companies, 1174 Marines in artillery batteries, 640 Marines in engineering companies, and 3071 Marines in the service support companies.

The data indicating service in the Gulf War, demographic variables for Gulf War troops, and military separation dates for Gulf War veterans were also obtained from DMDC. Because the comparison group (Marines returning from Vietnam) was exclusively male, and the small number of non-enlisted personnel had been excluded, only enlisted male Marines were included in the Gulf War study population as well. Similarly, because only 1st Marine Division troops were active at the end of the Vietnam conflict, the Gulf War study population was restricted to troops of the 1st Marine Division. An historical monograph²⁵ of 1st Marine Division activities during Desert Storm was used to verify the presence on the battlefield of the combat units used in this study.

The Gulf War study population was limited to the same four troop categories present at the end of USMC involvement in the Vietnam conflict: infantry troops, artillery troops, combat engineers, and service support troops. Seven infantry battalions of the 1st Marine Division were identified from the monograph as having actively participated in the ground war. The units composing the infantry group were the 1st, 2nd, and 3rd Battalions of the 7th Marine Regiment; the 1st Battalion of the 1st Marine Regiment; the 1st Battalion of the 5th Marine Regiment; the 3rd Battalion of the 9th Marines; and the Detachment 3rd Light Armored Infantry Battalion. Similarly, the 1st, 3rd, and 5th Battalions of the 11th Marines were identified in the monograph as active in the ground war and formed the artillery group for the Gulf War population. The 1st Combat Engineer Battalion was identified as actively participating during the ground war and constituted the group for the Gulf War that paralleled the combat engineer unit in the earlier conflict. Lastly, the service support group for the Gulf War was composed of troops from the following Force Service Support Group (FSSG) units: the Headquarters and Service Battalion, the 1st Supply Battalion, the 1st Maintenance Battalion, the 7th Motor Transport Battalion, the 1st Landing Support Battalion, the 7th Engineer Support Battalion, the 1st Medical Battalion, Marine Service Support Group – 11, Marine Service Support Group – 13, and the Headquarters Service Detachment. These service support units were identified as having been in the theater and having strengths greater than 100 personnel.

Altogether then, there were 10,878 Marines who were identified as having served in the Gulf in 1st Marine Division units that paralleled the units that returned from Vietnam at the end of that conflict. These 10,878 men were composed of 4374 Marines in infantry companies, 1588 Marines in artillery batteries, 616 Marines in engineering companies, and 4300 Marines in service support companies. Table 1 contrasts the numbers and types of troops at the beginning of the Vietnam and Gulf War postdeployment periods.

Data Sources – Hospitalizations

Hospitalization data for enlisted Marines returning from the Vietnam conflict were extracted from inpatient records maintained at the Naval Health Research Center. Diagnoses accompanying the hospital admissions incurred by Vietnam veterans during the five-year postdeployment period of this study were recorded in *International Classification of Diseases, Eighth Revision* (ICD-8) format.

Hospitalization data for enlisted Marines returning from the Gulf War were obtained from DMDC. Diagnoses accompanying the hospital admissions incurred by Gulf War veterans during the five-year postdeployment period of this study were recorded in *International Classification of Diseases, Ninth Revision* (ICD-9) format.

Each hospitalization record could contain up to eight diagnoses for a single admission. Hospitalization admissions for each deployed individual from the time of his return from the combat theater until the date five years from that return, or until his date of exit from service if less than five years, were included in the analyses. Individuals with injury admissions coded as resulting from battle wounds were excluded (DoD injury cause codes 400-479) from this study. Because reporting of admissions to the treatment facilities aboard aircraft carriers ceased in 1990, the six admissions aboard carriers recorded for the Vietnam cohort were removed from the analyses.

To eliminate re-counts of admissions that might be viewed as the same illness incident, the following criteria were used to aggregate admission records: any two disease admissions or any two injury admissions occurring in the same 7-day period were considered a single admission; disease admissions with identical 3-digit ICD diagnoses and occurring within the same 21-day period were counted as a single admission, as were follow-up admissions within this time period with the diagnosis 'no need for further medical care'; injury admissions with identical 3-digit ICD diagnoses and occurring within the same 60-day period were counted as a single admission. These adjustments

resulted in a 3.7% reduction in the total admission records for the Vietnam cohort and a 3.5% reduction in records for the Gulf War cohort. The mean time intervals between the aggregated records were 6.3 and 9.2 days respectively. In addition to eliminating re-counts of the same medical incident, the adjustment also ensured that the postdeployment admissions truly were post-deployment problems and not continuing treatments for a disease or injury sustained during the deployment.

STATISTICAL ANALYSES

Crude rates of hospitalization during the postdeployment periods were calculated for each of the four troop categories. The rates were calculated per 1000 persons per day (person-days). An individual who remained in the service for more than 5 years after his return from the conflict would contribute 1825 person-days (365×5) to the denominator (exposure period); an individual leaving the Marines exactly four months after his return from combat would contribute 120 person-days to the denominator. In addition to overall rates for the 5-year postdeployment period, rates were computed individually for each postdeployment year.

As a preliminary analysis indicated that there was a statistically significant difference in the ages of the returning veterans (mean age at return for Vietnam veterans of 22.3 versus 24.5 for Gulf War veterans), the crude hospital rates were age-adjusted. Because of the relative youth of forces deployed to the military engagements, age stratification for the adjustments was by narrow, four-year intervals. Table 2 shows the frequencies and percentages of person-days in each age group across the postdeployment periods. Age-adjustment was performed with the direct adjustment method, using the combined Vietnam and Gulf War group as the standardized population.²⁶ Ninety-nine percent confidence limits were computed for the age-adjusted rates.

Hospital admissions of the combat veterans groups were also compared by their proportional distribution within the following diagnostic categories: infective and parasitic, neoplasms, endocrine, blood and blood-forming, mental, nervous system, circulatory, respiratory, digestive, genitourinary, skin and subcutaneous tissue,

musculoskeletal, congenital anomalies, symptoms and ill-defined, injuries, adverse effects, supplementary classifications, and multiple categories. All categories except Injuries, Adverse Effects, and Supplemental Classifications directly correspond to the ICD diagnostic categories. However, in order to compare the proportions of admissions corresponding to 'injuries' alone, the diagnoses in the ICD category of 'Injuries and Poisonings' for adverse effects of medicinals and other external agents (ICD-9 diagnoses 960 – 994, 909) were separated into their own unique category. Additionally, admissions in the 'Injury and Poisonings' category coded as complications of surgical care (996-999) were placed with the Supplemental Classifications category admissions. These two minor data adjustments allowed 'injuries' to be reported as a stand-alone category of hospital admissions. All hospitalizations having diagnoses that fell into two separate diagnostic categories were reported in the 'multiple category' percentage. After the proportions corresponding to each diagnostic category were determined, 99% confidence limits were calculated for each category percentage.²⁷

RESULTS

Rate Comparisons

Figure 1 is a display of the age-adjusted hospitalization rates and confidence intervals of the four troop types composing the Vietnam and Gulf War veteran cohorts. For the Vietnam veterans, while there were no significant differences among the Artillery, Engineer, and Service Support groups, the rate of postdeployment hospitalizations for the Infantry troops was significantly higher when compared with those other three groups. Among Gulf War veterans, there were no statistically significant differences between any of the four troop categories.

Crude and age-adjusted hospitalization rates by post-conflict year for the Vietnam and Gulf War cohorts are presented in Table 3. Hospital admission rates for Vietnam veterans were significantly higher overall and for post-conflict years 1, 3, 4, and 5 when compared with the rates of Gulf War veterans. As can be seen in this table, there was a general decline in hospitalization rates among both groups in the years following the

conflicts. Of the veterans hospitalized, the Vietnam cohort and the Gulf War cohort both averaged 1.24 admissions.

Table 4 is a year-by-year comparison of postdeployment hospitalization rates for Infantry troops from the two conflicts. The rates of the Vietnam infantry troops were significantly higher than the Gulf War infantry troops across the five-year postdeployment period and in the first, fourth, and fifth post-conflict years. Table 5 is a display of the year-by-year comparisons between the Artillery troops serving in the Vietnam and Gulf War conflicts. There were no significant differences in postdeployment hospitalization rates among the artillery groups. Table 6 is a presentation of the yearly rate comparisons between the Combat Engineer groups serving in the two conflicts; there were no statistically significant differences between the hospitalization incidence incurred by the engineering groups active in the two conflicts. Table 7 contrasts the yearly postdeployment rates of the fourth study group -- Service Support troops. As can be seen in this table, the Service Support troops deployed to Vietnam exhibited a significantly higher rate of hospitalizations than Gulf War Service Support troops across the five-year post-conflict period, as well as significantly higher rates in the third and fifth postdeployment years.

Admission Type Comparisons

Table 8 contrasts the distribution of illness category types incurred by the Vietnam and Gulf War veterans across the five-year postdeployment period. Significant differences existed between the two groups of veterans in the proportions of 'Infective and Parasitic' diseases, 'Genitourinary' disorders, and 'Musculoskeletal' disorders, with the Vietnam veterans having the higher proportion within the first two categories and the Gulf War veterans having the higher proportion of musculoskeletal disorders. The specific three-digit ICD-9 diagnoses that were most prevalent among Gulf War veterans within the musculoskeletal category included: internal derangement of joint (30%), synovium tendon and bursa (16%), and other derangement of joint (14%). While slight differences exist in the coding between ICD-9 and ICD-8, the diagnoses of Gulf War veterans are

very similar to the types of musculoskeletal disorders incurred by Vietnam veterans: derangement of joint (25%), other unspecified disorder of joint (15%), synovitis bursitis tenosynovitis (12%), and intervertebral disc disorder (10%).

The difference in proportions of infective disorders was largely due to the considerably higher percentage of this category among Vietnam veterans in the very first year after their return from the combat theater. (Table 9). Similarly, it can be seen in Table 10 that the high percentages of infective disorders among the Vietnam veterans was largely due to those incurred by infantry and artillery troops. Because the percentages in Tables 9 and 10 are based on a large number of cells with frequencies of less than ten (50% and 48% respectively), confidence intervals were not computed for the proportions in these tables.

DISCUSSION

The present study sought to investigate any differences or similarities between the post-war hospitalization experiences of Gulf War and Vietnam veterans. Much controversy has surrounded the postdeployment illness incidence of Gulf War veterans, and the current study juxtaposed the hospital admission incidence of Marines deployed to the Gulf with the hospital incidence of a similar group of Marines deployed to an earlier military conflict. The strength of this study lies in the similarities of the comparison groups – hospitalization incidence of infantry, artillery, combat engineer, and service support units of the 1st Marine Division were contrasted for the years immediately following the veterans' return from two different conflicts. Similarly, both the Vietnam and Gulf War study groups were composed of troops that were in-theater at the end of USMC involvement in those conflicts. One caveat warrants mention here. The current investigation was only able to track the hospitalizations of Marines that remain in the service. Therefore, while the authors know of no evidence that suggests differences in hospitalization rates between veterans who remain in service and those that do not, the present study would not be able to track any such differences if they did exist.

Also, factors which might have been adjusted for if more recent data alone had been utilized, could not be adjusted for because of data unavailability within the Vietnam cohort; race and marital status are two such factors. Further, hospitalization rate differences could conceivably exist because of differences between the early 1970s and the early 1990s in preventive medicine techniques or in 'tendency to hospitalize' for certain conditions; those factors are likewise beyond the scope of this study. Nevertheless, the reported overall hospitalization rates of the Gulf War and Vietnam veterans in this study are based on 9.6 million and 6.6 million postdeployment person-days respectively, and therefore provide considerable insights into the post-conflict experiences of two groups of veterans.

Interestingly, while the infantry troops deployed to Vietnam showed the highest rate of hospital admissions among the four unit-types deployed to that conflict, there were no significant differences in hospitalization rates among the infantry, artillery, engineering, and service support units deployed to the Gulf War. Any hypothesized environmental exposure thought to have had a negative health effect on Gulf War veterans either affected the four troop types in a uniform fashion, or is not readily apparent through analyses of USMC hospitalizations.

Overall, and in four of the five postdeployment years, the aggregated Gulf War veterans exhibited lower hospitalization rates than did their Vietnam counterparts. Separately, the infantry and service support units deployed to the Persian Gulf had lower postdeployment hospitalization rates than did the Vietnam veterans. The combat engineer and artillery units deployed to the Gulf and Vietnam did not differ significantly in their respective rates of hospitalization. Also, among veterans who required hospitalization, the two cohorts did not differ in subsequent admissions; both the Vietnam and Gulf War groups averaged 1.2 hospital admissions among those who had at least one hospitalization.

The categories of illnesses incurred by the Gulf War veterans and the Vietnam veterans of this study were, by and large, not that dissimilar. The returning Vietnam veterans had

a higher percentage of their admissions comprised by infective/parasitic disorders and, to a lesser extent, genitourinary problems. The higher percentage of infectious disorders was largely due to the admissions in this category in the first year upon the veterans' return – a finding not unexpected given the tropical nature of the South East Asia theater of operations.²⁸ The Gulf War veterans had a higher proportion of their overall admissions comprised by musculoskeletal problems. However, the specific types of musculoskeletal disorders incurred by Gulf War veterans were not substantially different from their Vietnam counterparts. The prevalence of musculoskeletal complaints by Gulf War veterans in the DoD's Clinical Evaluation Program has previously been documented.²⁹

SUMMARY

The overall hospital admission rate of the Gulf War veterans of this study was significantly lower than the rate of similar troops deployed to the last ground conflict with sustained US involvement. Examined individually, the infantry and service support units deployed to the Gulf exhibited significantly lower hospitalization rates following the conflict than did their Vietnam counterparts; the rates of the engineer and artillery units deployed to the Gulf War did not differ significantly from similar units deployed to Vietnam. Among Gulf War-deployed Marines, the musculoskeletal disorder category represented a significantly higher proportion of their overall admissions than observed among the Vietnam veterans. In contrast, Vietnam veterans had significantly higher percentages of their overall admissions accounted for by infective/parasitic disorders and genitourinary disorders. With the exception of these categories of hospital admissions, illness category percentages between the Gulf War and Vietnam veterans exhibited but minimal differences. Among individuals who were hospitalized, the average number of admissions was the same for the Gulf War and Vietnam cohorts.

The lack of differences between the hospitalization rates of various Gulf War units (infantry, artillery, engineer, service support) performing different functions in varying locations on the battlefield, undercuts the notion that a specific environmental exposure is

responsible for deleterious health effects on Gulf War veterans; any specific risk factor might have been expected to affect these very different groups disproportionately. Indeed, Marines deployed to the Gulf War theater exhibited rates of hospitalization that are lower or comparable to the rates incurred by Marines returning from a previous ground combat operation. Nevertheless, the potential psychological disruption associated with deploying a young man or woman across the globe to risk his or her life defending national or international interests, under sometimes harrowing battlefield conditions, is not insubstantial. All appropriate measures should be taken to adequately prepare such individuals before deployment, to monitor their mental and physical well-being during the deployment, and to provide any appropriate counseling or treatment following their return from war.³⁰

REFERENCES

1. Haley RW, Kurt TL. Self-reported exposure to neurotoxic chemical combinations in the Gulf War: a cross-sectional epidemiologic survey. *JAMA*. 1997;277(3):231-237.
3. National Institutes of Health Technology Assessment Workshop Panel. The Persian Gulf experience and health. *JAMA*. 1994;272:391-396.
4. The Institute of Medicine. *Health Consequences of Service During the Persian Gulf War: Initial Findings and Recommendations for Immediate Action*. Washington, DC: National Academy Press; 1995.
4. The Institute of Medicine. *Health Consequences of Service During the Persian Gulf War: Recommendations for Research and Information Systems*. Washington, DC: National Academy Press; 1996.
5. Defense Science Board. *Report of the Defense Science Board Task Force on Persian Gulf War Health Effects*. Washington, DC: Office of the Under Secretary of Defense for Acquisition and Technology; 1994.
6. Presidential Advisory Committee on Gulf War Veterans Illnesses. *Final Report*. Washington, DC: U.S. Government Printing Office; December 1996.
7. Shenon P. Many veterans of the Gulf War detail illnesses from chemicals. *New York Times*. September 20, 1996: A1.
8. Haley RW, Kurt TL, Hom J. Is there a Gulf War syndrome?: searching for syndromes by factor analysis of symptoms. *JAMA*. 1997;277:215-222.
9. Haley RW, Billecke S, La Du BN. Association of PON1 type Q arylesterase activity with neurological symptom complexes in Gulf War Veterans. *Toxicol Appl Pharmacol*. 1999; 157(3):227-233.
10. Anger WK, Storzbach D, Binder LM, et al. Neurobehavioral deficits in Persian Gulf veterans: evidence from a population-based study. *J Int Neuropsychol Soc*. 1999;5(3):203-212.
11. Pennisi E. Chemicals behind Gulf War syndrome? *Science*. 1996;272(5261):479-480.
12. Petrucelli BP, Goldenbaum M, Scott, B, et al. Health effects of the 1991 Kuwait oil fires: a survey of US Army troops. *J Occup Environ Med*. 1999;41(6):433-439.
13. Durakovic A. Medical effects of internal contamination with uranium. *Croatian Medical Journal*. 1999;40(1):49-66.
14. Cook JE, Wenger CB, Kolka MA. Chronic pyridostigmine bromide administration: side effects among soldiers working in a desert environment. *Mil Med*. 1992;157:250-254.

15. Magill AJ, Grogl M, Johnson SC, Gasser RA Jr. Visceral infection due to *Leishmania tropica* in a veteran of Operation Desert Storm who presented 2 years after leaving Saudi Arabia. *Clin Infect Dis*. 1994;19:805-806.
16. Abou-Donia MB, Wilmarth KR, Jensen KF, Oehme FW, Kurt TL. Neurotoxicity resulting from coexposure to pyridostigmine bromide, DEET, and permethrin: implications of Gulf War chemical exposures. *J Toxicol Environ Health*. 1996;48:35-56.
17. Miller CS, Prihoda TJ. A controlled comparison of symptoms and chemical intolerances reported by Gulf War veterans, implant recipients and persons with multiple chemical sensitivity. *Toxicol Ind Health*. 1999;15(3-4),386-397.
18. Adler AB, Vaitkus MA, Martin JA. Combat exposure and posttraumatic stress symptomatology among US soldiers deployed to the Gulf War. *Military Psychology*. 1993;8(1):1-14.
19. Baker DG, Mendenhall CL, Simbartl LA, Magan LK, Steinberg JL. Relationship between posttraumatic stress disorder and self-reported physical symptoms in Persian Gulf War veterans. *Arch Intern Med* 1997;157:2076-2078.
20. Stretch RH, Marlowe DH, Wright KM, Bliese PD, Knudson KH, Hoover CH. Posttraumatic stress disorder symptoms among Gulf War veterans. *Mil Med*. 1996;161:407-410.
21. Gray GC, Coate BD, Anderson CM, et al. The postwar hospitalization experience of US veterans of the Persian Gulf War. *N Engl J Med*. 1996;335:1505-1513.
22. Kang HK, Bullman T. Mortality among US veterans of the Persian Gulf War. *N Engl J Med*. 1996;335(20):1498-1504.
23. Hyams KC, Wignall FS, Roswell R. War syndromes and their evaluation: from the U.S. Civil War to the Persian Gulf War. *Ann Intern Med*. 1996;125:398-405.
24. Stanton SL. *Vietnam Order of Battle*. Washington, DC: U.S. News Books; 1981.
25. Cureton CH. *U.S. Marines in the Persian Gulf, 1990-1991, With the 1st Marine Division in Desert Shield and Desert Storm*, History and Museums Division, Headquarters, U.S. Marine Corps, Washington, D.C., 1993.
26. Lilienfeld AM, Lilienfeld DE. *Foundations of Epidemiology*. New York, NY: Oxford University Press; 1980.
27. Zar JH. *Biostatistical Analysis*. Englewood Cliffs, NJ: Prentice-Hall, Inc.; 1984.
28. US Department of Health and Human Services. *Health Status of Vietnam Veterans, Volume III. Medical Examination*. Atlanta, GA: US Department of Health and Human Services; 1989.

29. Erickson AR, Enzenauer RJ, Bray VJ, West SG. Musculoskeletal complaints in Persian Gulf veterans. *J Clin Rheumatol*. 1998;4:181-185.
30. National Science and Technology Council. *Planning for Health Preparedness for and Readjustment of the Military, Veterans, and Their Families after Future Deployments*. Washington, DC: Executive Office of the President of the United States; 1998.

Table 1. Study Population by Troop Type and Cohort

USMC Troop Type	Vietnam	Gulf War
Infantry	7009	4374
Artillery	1174	1588
Engineering	640	616
Service Support	3071	4300
Total	11894	10878

Table 2. Person-days of postdeployment periods by Age-group and Cohort

Age (years)	Vietnam Veterans			Gulf War Veterans	
	Number of Person-days	Percent of Person-days		Number of Person-days	Percent of Person-days
18 - 21	2849217	43%		1822510	19%
22 - 25	1869323	28%		3665567	38%
26 - 29	603554	9%		1752292	18%
30 - 33	477639	7%		1093720	11%
34 - 37	465913	7%		717147	7%
38 - 41	207208	3%		383125	4%
42 - 45	95320	1%		153450	2%
46 - 49	34788	1%		46189	0%
> 50	14264	0%		12614	0%
All ages	6617226	100%		9646614	100%

Table 3 - Crude and Age-Adjusted Hospitalization Rates by Post-conflict Year and Cohort

Period of Observation	Vietnam Veterans		Gulf War Veterans	
	Crude Rate*	Age-Adjusted* (99%CI)	Crude Rate	Age-Adjusted (99% CI)
Year 1	0.327	0.312 (0.295, 0.330)†	0.230	0.233 (0.218, 0.249)
Year 2	0.250	0.243 (0.222, 0.263)	0.203	0.209 (0.190, 0.228)
Year 3	0.270	0.268 (0.241, 0.295)†	0.205	0.209 (0.186, .233)
Year 4	0.222	0.218 (0.190, 0.247)†	0.161	0.160 (0.136, 0.185)
Year 5	0.248	0.245 (0.212, 0.277)†	0.115	0.114 (0.092, 0.136)
Overall	0.286	0.272 (0.261, 0.282)†	0.199	0.206 (0.197, 0.215)

*Rates are computed per 1000 person-days.

† Rate is significantly higher than that of Gulf War cohort.

**Table 4. Crude and Age-adjusted Hospitalization Rates
for Infantry Units by Cohort and Post-conflict Year**

	Time Period	Crude Rate*	Age-Adjusted Rate*	99% Confidence Interval	
				Lower	Upper
Vietnam Veterans	Year 1	0.382	0.371†	0.343	0.399
	Year 2	0.260	0.255	0.224	0.286
	Year 3	0.251	0.248	0.209	0.287
	Year 4	0.253	0.249†	0.201	0.298
	Year 5	0.274	0.273†	0.218	0.328
Gulf War Veterans	Overall	0.318	0.305†	0.289	0.322
	Year 1	0.241	0.243	0.220	0.265
	Year 2	0.193	0.201	0.174	0.229
	Year 3	0.229	0.239	0.200	0.277
	Year 4	0.144	0.147	0.110	0.185
	Year 5	0.140	0.134	0.096	0.173
	Overall	0.207	0.216	0.202	0.230

*Rates are computed per 1000 person-days.

† Rate is significantly higher than that of Gulf War cohort.

**Table 5. Crude and Age-adjusted Hospitalization Rates
for Artillery Units by Cohort and Post-conflict Year**

	Time Period	Crude Rate*	Age-Adjusted Rate*	99% Confidence Interval	
				Lower	Upper
Vietnam Veterans	Year 1	0.238	0.235	0.191	0.279
	Year 2	0.181	0.184	0.135	0.232
	Year 3	0.216	0.221	0.153	0.288
	Year 4	0.188	0.167	0.097	0.237
	Year 5	0.163	0.139	0.070	0.207
	Overall	0.211	0.203	0.178	0.229
Gulf War Veterans	Year 1	0.237	0.240	0.195	0.284
	Year 2	0.275	0.277	0.217	0.338
	Year 3	0.186	0.182	0.121	0.243
	Year 4	0.185	0.187	0.113	0.261
	Year 5	0.086	0.085	0.031	0.139
	Overall	0.217	0.220	0.194	0.247

*Rates are computed per 1000 person-days.

† Rate is significantly higher than that of Gulf War cohort.

**Table 6. Crude and Age-adjusted Hospitalization Rates
for Combat Engineer Units by Cohort and Post-conflict Year**

	Time Period	Crude Rate*	Age-Adjusted Rate*	99% Confidence Interval	
				Lower	Upper
Vietnam Veterans	Year 1	0.240	0.249	0.181	0.317
	Year 2	0.307	0.301	0.203	0.400
	Year 3	0.228	0.220	0.114	0.327
	Year 4	0.166	0.170	0.063	0.277
	Year 5	0.198	0.265	0.119	0.411
	Overall	0.242	0.242	0.199	0.285
Gulf War Veterans	Year 1	0.191	0.209	0.146	0.271
	Year 2	0.252	0.255	0.164	0.346
	Year 3	0.221	0.260	0.144	0.375
	Year 4	0.175	0.172	0.064	0.279
	Year 5	0.057	0.053	0.000	0.119
	Overall	0.195	0.209	0.169	0.248

*Rates are computed per 1000 person-days.

† Rate is significantly higher than that of Gulf War cohort.

**Table 7. Crude and Age-adjusted Hospitalization Rates
for Service Support Units by Cohort and Post-conflict Year**

	Time Period	Crude Rate*	Age-Adjusted Rate*	99% Confidence Interval	
				Lower	Upper
Vietnam Veterans	Year 1	0.263	0.240	0.213	0.267
	Year 2	0.247	0.233	0.200	0.266
	Year 3	0.327	0.327†	0.280	0.375
	Year 4	0.196	0.195	0.153	0.237
	Year 5	0.244	0.240†	0.191	0.290
Gulf War Veterans	Overall	0.260	0.246†	0.229	0.262
	Year 1	0.223	0.224	0.198	0.250
	Year 2	0.179	0.182	0.153	0.212
	Year 3	0.188	0.189	0.153	0.226
	Year 4	0.164	0.165	0.127	0.204
	Year 5	0.112	0.113	0.079	0.147
	Overall	0.185	0.189	0.175	0.204

*Rates are computed per 1000 person-days.

† Rate is significantly higher than that of Gulf War cohort.

Table 8. Proportional Distribution of Hospitalizations by Diagnostic Category and Cohort

Diagnostic Category	Vietnam		Gulf War	
	Percent of admissions	99% CI	Percent of admissions	99% CI
Infectious & Parasitic Diseases	10.84	(9.05, 12.78)*	2.82	(1.91, 3.97)
Neoplasms	1.06	(0.55, 1.82)	1.25	(0.68, 2.06)
Endocrine, Metabolic & Immunity	0.53	(0.20, 1.13)	0.47	(0.16, 1.04)
Blood & Blood Forming Organs	0.26	(0.06, 0.74)	0.21	(0.04, 0.65)
Mental Disorders	10.20	(8.45, 12.05)	10.22	(8.35, 11.72)
Nervous System & Sense Organs	2.54	(1.69, 3.61)	1.88	(1.17, 2.85)
Circulatory System	3.12	(2.17, 4.29)	2.09	(1.35, 3.07)
Respiratory System	6.03	(4.70, 7.54)	6.78	(5.26, 8.27)
Digestive System	6.92	(5.49, 8.50)	9.65	(7.87, 11.16)
Genitourinary System	6.45	(5.07, 7.99)*	3.60	(2.54, 4.80)
Skin & Subcutaneous System	3.81	(2.75, 5.06)	2.50	(1.66, 3.61)
Musculoskeletal	9.36	(7.68, 11.16)*	21.07	(19.24, 22.59)
Congenital Anomalies	0.48	(0.17, 1.05)	0.57	(0.23, 1.18)
Symptoms, Signs, & Ill-Defined	2.54	(1.69, 3.61)	1.41	(0.79, 2.30)
Injuries	14.53	(12.50, 16.65)	12.42	(10.56, 13.96)
Adverse Effects	3.17	(0.23, 1.20)	0.52	(0.19, 1.11)
Supplemental	0.58	(2.22, 4.33)	3.03	(2.05, 4.25)
Multiple Categories	17.60	(15.32, 19.91)	19.51	(17.65, 20.94)

*Significant difference between Vietnam veterans and Gulf War Veterans diagnostic category proportions.


Table 9. Diagnosis Category Percentages by Year Across the Post-Deployment Periods

Diagnosis Category	Year 1		Year 2		Year 3		Year 4		Year 5	
	Vietnam %	Gulf %	Vietnam %	Gulf %	Vietnam %	Gulf %	Vietnam %	Gulf %	Vietnam %	Gulf %
Infectious & Parasitic Diseases	15.25	2.88	7.35	2.46	5.42	4.24	5.80	2.17	2.31	0.86
Neoplasm	0.80	1.38	1.84	1.44	0.83	1.21	0.00	0.54	2.31	0.86
Endocrine, Metabolic & Immunity	0.40	0.63	0.00	0.41	0.42	0.30	2.17	0.54	1.54	0.00
Blood & Blood Forming Organs	0.20	0.38	0.26	0.00	0.42	0.30	0.72	0.00	0.00	0.00
Mental Disorders	10.87	10.50	10.50	11.29	10.00	10.91	5.80	8.15	9.23	5.17
Nervous System & Sense Organs	2.59	1.75	3.15	2.26	1.25	0.91	3.62	2.72	1.54	2.59
Circulatory System	2.99	1.88	2.36	1.03	2.50	3.64	7.25	2.17	3.08	3.45
Respiratory System	5.28	7.88	7.87	6.37	5.83	7.27	7.25	4.89	5.38	2.59
Digestive System	6.18	9.63	6.56	9.45	8.33	8.48	6.52	10.87	11.54	12.07
Genitourinary System	6.68	3.50	3.94	2.26	8.75	4.85	7.25	3.26	6.92	6.90
Skin & Subcutaneous System	3.89	3.00	3.15	2.46	5.83	2.42	3.62	2.17	1.54	0.00
Musculoskeletal	6.98	17.25	8.92	20.12	13.33	24.85	14.49	24.46	16.15	35.34
Congenital Anomalies	0.30	0.63	1.05	0.82	0.00	0.61	0.72	0.00	0.77	0.00
Symptoms, Signs, & Ill-Defined	1.60	1.13	2.89	1.64	4.58	1.21	5.07	2.17	2.31	1.72
Injuries	13.36	15.75	17.59	13.96	16.67	6.97	11.59	10.33	13.85	1.72
Adverse Effects	0.30	0.38	0.79	1.23	0.00	0.30	1.45	0.00	2.31	0.00
Supplemental	3.59	3.00	3.15	3.70	1.67	3.03	2.90	1.63	3.08	2.59
Multiple Categories	18.74	18.50	18.64	19.10	14.17	18.48	13.77	23.91	16.15	24.14

Table 10. Admission Diagnosis Category Percentages by Unit Types

Diagnosis Category	Infantry		Artillery		Engineering		Service Support	
	Vietnam %	Gulf %	Vietnam %	Gulf %	Vietnam %	Gulf %	Vietnam %	Gulf %
Infectious & Parasitic Diseases	13.03	3.83	10.64	1.98	1.16	1.98	7.51	2.25
Neoplasms	0.69	1.45	2.13	0.66	2.33	0.00	1.38	1.46
Endocrine, Metabolic & Immunity	0.17	0.40	2.13	0.00	0.00	0.99	0.99	0.66
Blood & Blood Forming Organs	0.26	0.26	0.00	0.00	0.00	0.99	0.40	0.13
Mental Disorders	10.70	10.69	6.38	10.56	10.47	12.87	10.08	9.27
Nervous System & Sense Organs	2.59	1.72	4.26	1.65	1.16	2.97	2.17	1.99
Circulatory System	3.11	1.58	3.55	0.99	2.33	3.96	3.16	2.78
Respiratory System	5.61	6.60	9.93	8.25	6.98	8.91	5.73	6.09
Digestive System	6.13	9.10	9.22	9.90	8.14	6.93	7.91	10.46
Genitourinary System	6.99	4.49	6.38	4.29	5.81	0.00	5.34	2.91
Skin & Subcutaneous System	3.54	2.37	2.84	2.31	8.14	2.97	3.95	2.65
Musculoskeletal	8.11	20.05	9.22	19.47	12.79	20.79	11.66	22.78
Congenital Anomalies	0.52	0.79	0.71	0.66	0.00	1.98	0.40	0.13
Symptoms, Signs, & Ill-Defined	2.67	1.45	0.00	1.32	3.49	0.00	2.77	1.59
Injuries	14.84	14.12	12.06	11.22	15.12	11.88	14.43	11.26
Adverse Effects	0.78	0.79	0.71	0.33	0.00	1.98	0.20	0.13
Supplemental	2.59	3.69	4.26	3.30	3.49	0.00	4.15	2.65
Multiple Categories	17.69	16.62	15.60	23.10	18.60	20.79	17.79	20.79

Figure 1. Age-Adjusted Postdeployment Hospitalization Rates by Troop Type



APPENDIX B

Christopher G. Blood
Naval Health Research Center
P.O. Box 85122
San Diego, CA 92186-5122
(619) 553-8386

PERSONAL QUALIFICATIONS AND RESEARCH CONTRIBUTIONS

A. ACADEMIC BACKGROUND

1. J.D. in Law from Thomas Jefferson School of Law; expected graduation Aug 2001.
2. M.A. in Research Psychology from San Francisco State University, May 1981.
3. B.A. from University of California, Santa Barbara in Experimental Psychology, June 1979.

B. PROFESSIONAL EXPERIENCE

1. Operations Research Division, Naval Health Research Center, 1993 through present.
2. Research Psychologist, Naval Health Research Center May 1982 through present.
3. Member of *Scientific Planning and Review Council*, Naval Health Research Center; Council is responsible for reviewing the scientific merit of all research proposals and charting the direction of the Center, 1994 - 1997.
4. *Contracting Officer's Technical Representative* for Contract N66001-93-D-0021 (a 12 million dollar biomedical research support contract), June 1992 - January 1996. Was responsible for writing Request for Proposals, Statement of Work, setting evaluation criteria, evaluating proposals, and administering contract as the technical representative of Navy Contracting.
5. Member of the *Committee for the Protection of Human Subjects*, April 1988 - August 1992; CPHS is responsible for ensuring the safety and informed consent of all study participants.

C. RESEARCH CONTRIBUTIONS

Journal Publications

Derderian BR, Blood CG. Shipboard Medical Admissions During Peacetime and Combat Support Deployments. Military Medicine, Vol. 165 (3), 228-36, March 2000.

Walker GJ, Blood CG. The Patient Flow of Marine Disease and Non-Battle Injury Conditions Within a Multi-Echelon System of Care. Military Medicine, Vol. 164, 423-27, June 1999

Walker GJ, Blood CG. The Patient Flow of Wounded Marines Within a Multi-Echelon System of Care. Military Medicine, Vol. 168, October 1999

Blood CG. Illness Incidence During Military Operations As A Soft Operations Factor, Military Operations Research, Volume 2 (3), December 1996.

Blood CG, Odowick MS, Marks JS. Simulation of Casualty Sustainment During Naval Combat Operations, Simulation, 67:6, 367-374, December, 1996.

Blood CG. Blue on Blue: A History of Friendly Fire, *Book Review*. Naval War College Review, Winter, 1997.

Blood CG, Jolly RT, Odowick MS. Casualty Incidence During Naval Combat Operations: A Matter of Medical Readiness, Naval War College Review, Autumn, 1996.

Blood CG, O'Donnell ER. A System to Project Injury and Illness Incidence During Military Operations, Journal of Medical Systems, 19, 6: 457-64, 1995.

Blood CG, Jolly RT. Comparisons of Disease and Nonbattle Injury Incidence Across Various Military Operations, Military Medicine, 160, 5: 258-63, 1995.

Gauker ED, Blood CG. Friendly Fire Incidents during World War II Naval Operations, Naval War College Review, XLVIII, 1: 115-21, 1995.

Blood CG, Gauker ED, Jolly RT, Pugh WM. Comparisons of Casualty Presentation and Admission Rates during Various Combat Operations, Military Medicine, 159, 6: 457-61, 1994.

Blood CG, Anderson ME. The Battle for Hue: Casualty and Disease Rates during Urban Warfare, Military Medicine, 159, 7: 1994.

Burkle FM, Newland C, Orebaugh S, Blood CG. Emergency Medicine in the Persian Gulf--Part 2: Triage Methodology and Lessons Learned, Annals of Emergency Medicine, 23, 4:748-54, 1994.

Burkle FM, Newland C, Meister SJ, Blood CG. Emergency Medicine in the Persian Gulf--Part 3: Battlefield Casualties. Annals of Emergency Medicine, 23, 4:755-60, 1994.

Blood CG, Anderson ME, Gauker ED. Casualty Rates Among Naval Forces Ashore. Military Medicine, 158, 12:774-77, 1993.

Leedham CS, Blood CG. A Descriptive Analysis of Wounds among U.S. Marines Treated at Second-Echelon Facilities in the Kuwaiti Theater of Operations, Military Medicine, 158, 8:508-12, 1993.

Blood CG, Gauker ED. The Relationship Between Battle Intensity and Disease Rates among Marine Corps Infantry Battalions. Military Medicine, 158, 5:340-44, 1993.

Blood CG, Pugh WM, Gauker ED, et al. Comparisons of Wartime and Peacetime Disease and Non-Battle Injury Rates Aboard Ships of the British Royal Navy. Military Medicine, 157, 12:641-4, 1992.

Blood CG. Analyses of Battle Casualties by Weapon Type Aboard U.S. Navy Warships. Military Medicine, 157, 3:124-30, 1992.

Blood CG, Nirona CB. Outpatient Illness Incidence Aboard U.S. Navy Ships During and Following the Vietnam Conflict. Military Medicine, 155, 10:472-476, 1990.

Blood CG, Griffith DK. Ship Size as a Factor in Illness Incidence Among U.S. Navy Vessels. Military Medicine, 155, 7:310-314, 1990.

Blood CG & Brand DMP. Shipboard Medical Information Systems: Familiarity and Receptivity. Journal of Medical Systems, 1988, 12(3), 147-152.

Hoiberg AL, Blood CG. Health Risks of diving among U.S. Navy Officers. Undersea Biomedical Research, 1986, 13(2), 237-245.

Blood CG, Hoiberg AL. Analyses of Variables Underlying U.S. Navy Diving Accidents. Undersea Biomedical Research, 1985, 12(3), 252-261.

Hoiberg AL & Blood CG. Age-specific Morbidity and Mortality Rates Among U.S. Navy Divers and Controls. Undersea Biomedical Research, 1985, 12(2), 191-203.

Hoiberg AL, Blood CG. Age-specific Morbidity Among Navy Pilots. Aviation, Space, and Environmental Medicine, 1983, 54, 912-918.

Conference Presentations

Blood CG. Chairperson of Battlefield Performance, Casualty Sustainment, and Medical Planning Working Group at 66th Military Operations Research Society Symposium held at the Naval Postgraduate School, Monterey, CA, 23-25 June, 1998.

Blood CG. Projections of Ground Casualties and Shipboard Casualties During Military Operations. Presented at 66th Military Operations Research Society Symposium held at the Naval Postgraduate School, Monterey, CA, 23-25 June, 1998.

Blood CG. Casualty Projections for Ground Forces. 65th Military Operations Research Society Symposium, Quantico, VA, 10 - 12 June 1997.

Blood CG. Casualty Estimation for Forces Afloat. 65th Military Operations Research Society Symposium, Quantico, VA, 10 - 12 June 1997.

Blood CG. Projecting Battlefield Casualties: Incorporating Socio-Cultural and Other Adversary-Specific Factors. 64th Military Operations Research Society Symposium, Fort Leavenworth, KS, 18 - 20 June 1996.

Blood CG, Sundstrom SC. The Optimal Placement of Casualty Evacuation Assets: A Linear Programming Model. 64th Military Operations Research Society Symposium, Fort Leavenworth, KS, 18 - 20 June 1996.

Blood CG. Simulation of Casualty and Illness Incidence Among Ground Troops and Forces Afloat. Presented at the Twelfth International Symposium on Military Operation Research (12 ISMOR) Royal Military College of Science, Swindon, U.K. 4-8 September 1995.

Blood CG. Using the FORECAS Projection System to Simulate Casualty and Illness Rates Among Ground Forces, 63rd Military Operations Research Society Symposium, Annapolis MD, 6-8 1995.

Blood CG. Using the SHIPCAS Projection System to Simulate Casualty and Illness Rates Among Ground Afloat, 63rd Military Operations Research Society Symposium, Annapolis MD, 6-8 June 1995.

Blood CG. The Battle for Hue: Casualty and Disease Rates during Urban Warfare. Presented at the Tenth International Symposium on Military Operation Research (10 ISMOR) Royal Military College of Science, Swindon, U.K. 7-10 September 1993.

Blood CG. Illness Incidence During Military Operations as a Soft OR Factor, 61st Military Operations Research Society Symposium, Dayton, Ohio, 22-24 June 1993.

Blood CG. The Relationship Between Battle Intensity and Disease Rates Among Combat and Support Units. Presented at the Ninth International Symposium on Military Operation Research (9 ISMOR) Royal Military College of Science, Swindon, U.K. 7-11 September 1992.

Technical Reports

Singh S, Blood CG, Zouris JM. Projection of Patient Condition Code Distributions for Naval Combat Deployments, Naval Health Research Center Report No. 99-20.: San Diego, CA, 1999.

Walker GJ, Zouris JM, Blood CG. Projection of Patient Condition Code Distributions During Ground Operations, Naval Health Research Center Report No. 99-17.: San Diego, CA, 1999.

Derderian BR, Blood CG. Shipboard Medical Admissions During Peacetime and Combat Support Deployments. Naval Health Research Center Report No. 98-30.: San Diego, CA, 1998.

Walker GJ, Blood CG. The Patient Flow of Marine Disease and Non-Battle Injury Conditions Within a Multi-Echelon System of Care. Naval Health Research Center Report No. 98-11.: San Diego, CA 1998.

Walker GJ, Blood CG. The Patient Flow of Wounded Marines Within a Multi-Echelon System of Care. Naval Health Research Center Report No. 98-8.: San Diego, CA, 1998.

Blood CG, Zouris JM, O'Donnell ER, Rotblatt D. Using the Ground Forces Casualty Forecasting System (FORECAS) to Project Casualty Sustainment, San Diego (CA): Naval Health Research Center; 1997, Report No. 97-39.

Matheny SA, Keith DC, Sundstrom SC, Blood CG. A Medical Planning Tool for Projecting Casualty Evacuation Assets in a Military Theater of Operations, San Diego (CA): Naval Health Research Center; 1997, Document No. 97-3.

Blood CG, Marks JS, Le LP. Using the Shipboard Casualty Projection System (SHIPCAS) to Forecast Ship Hits and Casualty Sustainment, San Diego (CA): Naval Health Research Center; 1997, Report No. 97-3.

Blood CG, Marks JS, Odowick MS. Shipboard Casualty Forecasting: Adjustments To Ship Hit Probabilities, San Diego (CA): Naval Health Research Center; San Diego (CA): Naval Health Research Center; 1996, Report No. 96-25.

Marks JS, Blood CG, Gilman PA. Casualty Sustainment During Naval Warfare: Adjustments to World War II-Based Projections, San Diego (CA): Naval Health Research Center; 1996, Report No. 95-37.

Blood CG, Jolly RT, Odowick MS. An Analysis of Shipboard Casualty During Naval Combat Operations, San Diego (CA): Naval Health Research Center; 1995, Report No. 95-13.

Odowick MS, Blood CG. Using the SHIPCAS Projection System to Simulate Casualty and Illness Rates Among Forces Afloat, San Diego (CA): Naval Health Research Center; 1994, Document No. 94-6F.

O'Donnell ER, Blood CG. A Simulation Program to Project Casualty and Illness Rates. San Diego (CA): Naval Health Research Center; 1994, Document No. 94-2B.

O'Donnell ER, Blood CG. A Bivariate AutoRegressive Integrated Moving Average Analysis of Combat Troop Casualty Rates, San Diego (CA): Naval Health Research Center; 1994, Rep. No. 94-2.

Gauker ED, Blood CG. Amicide at Sea: Friendly Fire Incidents during World War II Naval Operations, San Diego (CA): Naval Health Research Center; 1994, Report No. 94-1.

Blood CG, Gauker ED, Anderson ME, Odowick MS, O'Donnell ER. Projections of Battlefield Medical Casualties among U.S. Marine Forces for Various Theaters of Operations. 1993. San Diego.: (CA) Naval Health Research Center Report No. 93-35.

O'Donnell ER, Blood CG. Distribution Characteristics of Marine Corps Casualty and Illness Rates, San Diego (CA): Naval Health Research Center; 1993, Report No. 93-20.

Blood CG, Anderson ME. The Battle for Hue: Casualty and Disease Rates during Urban Warfare, San Diego.: Naval Health Research Center Report No. 93-16.

Blood CG, Gauker ED, Jolly R, Pugh WM. Comparisons of Medical Presentation and Admission Rates during Various Combat Operations, San Diego.: Naval Health Research Center Rep. No. 93-12.

Gauker ED, Blood CG. Severity of Battle Injuries Occurring Aboard U.S. Naval Warships. San Diego.: Naval Health Research Center. Report No. 92-11.

Leedham CS, Blood CG. A Descriptive Analysis of Wounds Among U.S. Marines Treated at Second Echelon Facilities in the Kuwaiti Theater of Operations. San Diego.: Naval Health Research Center. Report No. 92-6.

Blood CG, Anderson ME, Gauker ED. Casualty Rates among Naval Forces Ashore. San Diego.: Naval Health Research Center. Report No. 92-2.

Blood CG, Gauker ED. The Relationship Between Battle Intensity and Disease Rates among Marine Corps Infantry Battalions. San Diego.: Naval Health Research Center. Report 92-1.

Blood CG, Pugh WM. The Dynamics of Casualty Rate Estimation, Medical Resource Allocation, and Disease and Injury Surveillance. San Diego.: Naval Health Research Center. Report No. 91-15, 1991.

Blood CG, Pugh WM, Pearsall DM. Comparisons of Wartime and Peacetime Disease and Non-Battle Injury Rates Aboard Ships of the British Royal Navy. San Diego, Calif.: Naval Health Research Center. Report No. 91-2, 1991.

Blood CG. Analyses of Battle Casualties by Weapon Type Aboard U.S. Navy Warships. San Diego, Calif.: Naval Health Research Center. Report No. 91-1, 1991.

Blood, CG. Shipboard and Ground Troop Casualty Rates Among Navy and Marine Corps Personnel during World War II Operations. San Diego, Calif.: Naval Health Research Center, Report No. 90-16, 1990.

Blood CG, Nirona CB. Geographical and Temporal Variations in Outpatient Morbidity at U.S. Navy Overseas Facilities. San Diego, Calif.: Naval Health Research Center, Report No. 90-13, 1990.

Pugh WM, White MR, Blood CG. Disease and Non-battle Injury Rates for Enlisted Personnel During Peacetime. San Diego, Calif.: Naval Health Research Center, 1989, Report No. 89-51.

Blood CG, Nirona CB, Pederson LS. Medical Resource Planning: The Need to Use a Standardized Diagnostic System. San Diego, Calif.: Naval Health Research Center, 1989, Report No. 89-41.

Blood CG, Griffith DK, Nirona CB. Medical Resource Allocation: Injury and Disease Incidence Among Marines in Vietnam. San Diego, Calif.: Naval Health Research Center, 1989, Report No. 89-36.

Blood CG, Nirona CB. Outpatient Illness Incidence Aboard U.S. Navy Ships During and Following the Vietnam Conflict. San Diego, Calif.: Naval Health Research Center, 1989, Report No. 89-15.

Blood CG, Griffith DK. Ship Size as a Factor in Illness Incidence. San Diego, Calif.: Naval Health Research Center, 1988, Report No. 88-48.

Blood CG, Pugh WM, Griffith DK, Nirona CB. Navy Medical Resource Planning: Rates of Illness for Various Operational Theaters. San Diego, Calif.: Naval Health Research Center, 1988, Report No. 88-42.

Blood CG, Brand DMP, Pugh WM. Shipboard Medical Department Information Flows. San Diego, Calif.: Naval Health Research Center, 1987, Report No. 87-32.

Blood CG, Brand DMP. Shipboard Medical Information Systems: Familiarity and Receptivity. San Diego, Calif.: Naval Health Research Center, 1987, Report No. 87-31.

Blood CG, Peeps-Brand DM, Pugh WM, Helmkamp JC. Shipboard Medical Information System Needs Aboard Surface Ships. San Diego, Calif.: Naval Health Research Center, 1987, Report No. 87-18.

Helmkamp JC, Seidman JH, Bone CM, Blood CG, Kelley, JB. The Occurrence of Hearing Loss in a Cohort of Civilians Employed at a U.S. Navy Industrial Facility. San Diego Calif.: Naval Health Research Center, 1986, Report No. 86-34.

Hoiberg AL & Blood CG. Health Risks of Diving Among U.S. Navy Officers. San Diego, Calif.: Naval Health Research Center, 1985, Report No. 85-23.

Hoiberg AL & Blood CG. Age-specific Morbidity and Mortality Rates Among U.S. Navy Divers and Controls. San Diego, Calif.: Report No. 84-25.

Blood CG, Hoiberg A. Diving Accidents: Analyses of Underlying Variables. San Diego, Calif.: Naval Health Research Center, 1984, Report No. 84-14.

Hoiberg AL, Blood CG. Effects of Age and Exposure on the Health Status of U.S. Navy Divers. San Diego, Calif.: Naval Health Research Center, 1984, Report No. 84-3.

Hoiberg AL, Blood CG. Age-specific Morbidity Among Navy Pilots. San Diego, Calif.: Naval Health Research Center, 1982, Report No. 82-32.



DEPARTMENT OF THE ARMY
US ARMY MEDICAL RESEARCH AND MATERIEL COMMAND
504 SCOTT STREET
FORT DETRICK, MARYLAND 21702-5012

REPLY TO
ATTENTION OF:

MCMR-RMI-S (70-1y)

19 Jan 01

MEMORANDUM FOR Administrator, Defense Technical Information
Center, ATTN: DTIC-OCA, 8725 John J. Kingman
Road, Fort Belvoir, VA 22060-6218

SUBJECT: Request Change in Distribution Statement

1. The U.S. Army Medical Research and Materiel Command has reexamined the need for the limitation assigned to technical reports written for Grant DAMD17-98-1-8010. Request the limited distribution statement for Accession Document Number ADB254366 be changed to "Approved for public release; distribution unlimited." This report should be released to the National Technical Information Service.

2. Point of contact for this request is Ms. Judy Pawlus at DSN 343-7322 or by email at judy.pawlus@det.amedd.army.mil.

FOR THE COMMANDER:

PHYLLIS M. RINEHART
Deputy Chief of Staff for
Information Management